



9353 Hill Road • Swartz Creek, MI 48473
(810) 635-4400 • Fax (810) 635-4404

www.lakeproinc.com

2018 Island Lake Water Quality Review

Introduction

The goals of this testing protocol were to monitor various water quality parameters of the lake, compare results to historical data, and identify any potential risks to the health of Island Lake. Water samples were taken at two different locations and tested for 14 parameters. Tests were conducted on a monthly basis from April through August. This report describes conditions at the times the samples were taken. The quality of the water was tested only to the parameters listed below.

Test results were compared to historical data from the report “2017 Island Lake Water Quality Review” by LakePro, Inc.

In this report, we included historical data from Water Quality Investigators. Their report provided annual averages for many of the parameters from 2002 to 2009. Including this data allowed us to see more accurate trends in the water quality data. In order to make the analysis easier, we added annual averages for our data and trendlines on the graphs. The trend lines provide quick indication of how each water quality parameter changed over the testing history.

Results

Parameter	2018 Season		
	Average	Target Range	Status
Temperature	69.8 °F	Less Than 75 °F	● Healthy
Dissolved Oxygen	7.7 mg/L	4.0 – 12.0 mg/L	● Healthy
Total Phosphorus	106 ppb	0 – 100 ppb	● Slightly High
Phosphate	37 ppb	0 – 100 ppb	● Healthy
Nitrate	462 ppb	0 – 1,000 ppb	● Healthy
Chlorophyll-a	7.0 ppb	0 – 7.3 ppb	● Healthy
Transparency	7.8 feet	More than 6.5 feet	● Healthy
pH	8.1 S.U.	7.0 – 9.0 S.U.	● Healthy
Total Dissolved Solids	374 ppm	0 – 1,000 ppm	● Healthy
Conductivity	748 ppm	0 – 1,500 ppm	● Healthy
Alkalinity	132 ppm	100 – 250 ppm	● Healthy
Sulfate	13.2 ppm	3 – 30 ppm	● Healthy
Fluoride	0.09 ppm	0.01 – 0.30 ppm	● Healthy
Chloride	126 ppm	0 – 230 ppm	● Healthy



9353 Hill Road • Swartz Creek, MI 48473
(810) 635-4400 • Fax (810) 635-4404

www.lakeproinc.com

Year-End Discussion



Island Lake’s water quality was excellent this year. The season average for all parameters were within the target ranges except for Total Phosphorus, which was slightly above the target range.

Temperature and Dissolved Oxygen



The average **Temperature** of the surface water was within the target range, but the water temperatures rose above the limit in June and stayed through August. Colder water can hold more oxygen, so cooler water is preferred for healthy lake ecology. Despite the water temperatures, the **Dissolved Oxygen** concentrations stayed within the target range for the entire summer.

We also measured temperature at different depths to create a temperature profile. This data shows how the temperature changed with depth and whether or not a thermocline was present in the lake. The first graph below shows the data we collected in 2018. This summer, there was a thermocline in the water, between ten and twenty feet. In the spring, the water temperature decreased steadily from the surface to the lake bottom.

We used the water temperature and volume at each depth to calculate a volume weighted average. The results of those calculations are presented in the second graph below. The trend line shows that over our testing history, the average temperatures of the lake remained steady.



The depth profile methodology was repeated and measurements taken for dissolved oxygen. This allowed us to see how the oxygen concentrations changed throughout the water column. We also used the dissolved oxygen and volume at each depth to calculate a volume weighted average. The trend line shows that over our testing history, the dissolved oxygen of the entire lake increased.







9353 Hill Road • Swartz Creek, MI 48473
(810) 635-4400 • Fax (810) 635-4404

www.lakeproinc.com


Nutrients, Plant Production, and Transparency

Nutrients in the water are the fuel for plant growth. Measuring the nutrient concentrations reveals the potential for nuisance plant growth. Phosphorus is a major nutrient necessary for aquatic plant growth, so it is important that this nutrient remains low in the lake. The **Total Phosphorus** was elevated in April, May, and June then returned to the target range in late summer. **Phosphate**, which is the form of phosphorus usable to plants, was within the target range for all samples and tests.

The depth profile methodology was repeated and measurements taken for total phosphorus. This revealed how this nutrient concentration changed throughout the water column. We also used the concentration and volume at each depth to calculate a volume weighted average. The trend line shows that over our testing history, the total phosphorus within the entire lake increased.



Additionally, we used the concentration of phosphorus and the lake volume to calculate the total phosphorus load. This quantified the amount of phosphorus in the lake. The following graph shows the results of this calculation over the years of our testing. Our industry touts 1 pound of phosphorus can support 500 pounds of plant growth, so the average load of 658 pounds of phosphorus could support almost 165 tons of plants!



Nitrate is another major nutrient for aquatic plant growth. The nitrate concentrations remained within the target range this summer. Although concentrations were in the target range, it is important that residents take measures to ensure their properties are not contributing excess fertilizers to the lake.

We also measured **Chlorophyll** concentrations because it is a reliable indicator of plant production. The target for chlorophyll is below 7.3 parts per billion. The chlorophyll concentrations were below the target level to start the summer, but crept above the target range in June and remained elevated through August. The abundant plant growth and moderate algae growth correlated with these results.



9353 Hill Road • Swartz Creek, MI 48473
 (810) 635-4400 • Fax (810) 635-4404

www.lakeproinc.com

A major effect of plant growth on the lake is the reduction of water clarity. Before algae forms the green mats of “scum” on the surface, it is suspended in the water column. Algae floating in the water can decrease water clarity even before you see a tint of green. This year, the **Transparency** averaged a depth of 7.8 feet. The April measurement was low, but clarity was very good in May through July. Clearer water is generally a positive attribute, but it also allows more sunlight to reach the lake bottom to fuel plant growth.

Trophic State Indices

In order to better understand the relationship between nutrients, plant production, and clarity, limnologists use Trophic State Indices (TSI) to score each category and examine the relationship between them. In general, lower scores indicate a less productive lake. The TSIs for Island Lake this year were:

Category	Water Quality Parameter	Trophic State Index (season average)	Classification
Nutrients	Total Phosphorus	71	Hypereutrophic
Plant Production	Chlorophyll	49	Mesotrophic
Clarity	Transparency	49	Mesotrophic



The TSI for total phosphorus classified the lake as hypereutrophic, or extremely productive, based on the availability of nutrients to fuel plant growth. The TSI for chlorophyll was lower than the nutrient index. This shows that despite the availability of nutrients, the plants were not at the levels predicted by the nutrient concentrations. This was due, in part, to the plant management on the lake. Finally, the TSI for transparency matched the production index and classified the lake as mesotrophic. This shows that the water clarity was consistent with the measured chlorophyll concentrations. Furthermore, the clarity was not degraded by dissolved solids or suspended particles.

Water Chemistry Parameters

It is important to monitor the basic water chemistry of the lake water. Shifts in these parameters can indicate major changes to the lake that may need to be further investigated.

The **pH** of the lake remained within the target range across all tests this year. This showed that the pH did not fluctuate to a point of concern despite changes in dissolved oxygen, alkalinity, and rainfall.

The depth profile methodology was repeated and measurements were taken for pH. This revealed how this parameter changed throughout the water column. We also used the readings and volume at each depth to calculate a volume weighted average. The trend line shows that over our testing history, the pH of the entire lake decreased.





9353 Hill Road • Swartz Creek, MI 48473
(810) 635-4400 • Fax (810) 635-4404

www.lakeproinc.com

The **Total Dissolved Solids (TDS)** were in the bottom of the target range throughout summer, showing there were low amounts of dissolved molecules in the water. This parameter includes nutrients, salts, and other substances, so it is a positive that this parameter remained low.



Conductivity measures ionic molecules in the water and usually follows the TDS. This parameter measures the molecules in the water ability to conduct electricity. So, it is particularly sensitive to salts, which are excellent conductors. The conductivity was in the middle of the target range this year, indicating a normal amount of ionic molecules in the lake and no immediate concern of salts.

Alkalinity measures the concentration of one salt, Calcium Carbonate, which is beneficial to the aquatic ecosystem. The carbonate ions are able to accept protons from acids, making it a natural buffer. This means that as acidic substances enter the lake, the carbonate is able to buffer against severe changes in pH that would pose a threat to the ecosystem. The alkalinity was at a healthy concentration for all tests this year.


Pollutants

The lake was tested for **Sulfate**, **Fluoride**, and **Chloride** as indicators of pollution. These molecules should be present in the water naturally, but elevated levels can indicate pollution from within the watershed and may pose a risk to the ecosystem. All three parameters were within their target ranges for all tests.

The depth profile methodology was repeated and measurements were taken for chloride. This allowed us to see how this parameter changed throughout the water column. We also used the results and volume at each depth to calculate a volume weighted average. The trend line shows that over our testing history, the chloride in the lake decreased slightly.



Finally, we used the concentration of chloride and the water volume of the lake to calculate the lake's chloride load. The following graph shows the results of this calculation over the years of our testing. This quantifies the amount of chloride in the lake.






9353 Hill Road • Swartz Creek, MI 48473
 (810) 635-4400 • Fax (810) 635-4404

www.lakeproinc.com


Historical Trends



Lake Temperature (°F)					
	April	May	June	July	August
West	51.0	69.2	76.4	78.4	75.9
East	49.2	68.8	76.2	78.0	75.3
Season Average					69.8

Target Range: < 75°F

The long-term trend for water temperature was generally flat and the 2018 season average was right on track. Water temperature is dependent upon air temperatures and the dates selected for testing. For that reason, LakePro tried to select similar dates for testing each year. The primary concern with warmer water temperatures decreasing oxygen solubility.



Dissolved Oxygen (mg/L)					
	April	May	June	July	August
West	9.1	8.5	7.4	6.4	6.9
East	9.4	8.7	7.3	6.4	6.6
Season Average					7.7


Target Range: 4.0 – 12.0 mg/L

Dissolved oxygen concentrations varied but followed a flat trend that corresponds to steady temperatures. Although the concentrations varied through the testing history, there has always been adequate oxygen for a healthy aquatic ecosystem.



9353 Hill Road • Swartz Creek, MI 48473
 (810) 635-4400 • Fax (810) 635-4404

www.lakeproinc.com




Total Phosphorus (ppb)					
	April	May	June	July	August
West	140	120	100	90	90
East	120	100	110	90	100
Season Average					106
Trophic State Index					71

Target Range: 0 – 100 ppb

Lakes generally accumulate the substances and materials that flow into it, including nutrients. Since testing began in 2003, the trend for phosphorus in the surface water tracked upward. The 2008 and 2009 concentrations were significantly higher than other years of testing, which pulled the trendline up. Since 2010 total phosphorus steadily increased, slipping above the target range in 2018.

There are ways to remove phosphorus, such as mechanical harvesting and heavy rain events that create excess flow. However, it is much easier to prevent excess nutrients from entering the lake. For this reason, it is vital that residents around the lake fertilize and use their land responsibly to prevent phosphorus from reaching the lake.



Phosphate (ppb)					
	April	May	June	July	August
West	10	40	40	40	40
East	20	40	50	40	50
Season Average					37


Target Range: 0 – 100 ppb

Phosphate is one form of phosphorus that is usable by plants and algae. So, as the total phosphorus accumulates, so too will the phosphate. There is no historical data available for phosphates, so the trendline reflected the changes only during the seven years of LakePro’s testing. The trend was upward, but the phosphate concentrations remained within the target range.



9353 Hill Road • Swartz Creek, MI 48473
 (810) 635-4400 • Fax (810) 635-4404


www.lakeproinc.com



Nitrate (ppb)					
	April	May	June	July	August
West	352	440	528	484	440
East	308	528	484	528	528
Season Average					462

Target Range: 0 – 1,000 ppb

Nitrate is another major nutrient that accumulates in lakes. The historical trend for nitrate is downward, due to consistently lower values since 2012 and despite wild variability in previous years of testing. It is important that residents fertilize and use their land responsibly to prevent additional nitrates from reaching the lake.



Chlorophyll (ppb)					
	April	May	June	July	August
West	3.7	5.0	8.4	9.2	7.4
East	4.1	5.7	10.4	8.1	7.9
Season Average					7.0
Trophic State Index					49

Target Range: 0 – 7.3 ppb


The linear trend for Chlorophyll was slightly downward. However, the average concentrations show a strong cyclical pattern. Except for 2008, the chlorophyll remained within the target range. Continuing the plant management activities will continue to suppress the algae and plants to acceptable levels. Continuing to mechanically harvest aquatic plants and to conduct herbicide treatments for dense invasive species will help keep the plant community to acceptable levels.

Another key to limiting plant production is to slow the influx of nutrients from both around the lake and throughout the watershed. Responsible land use practices, especially fertilizing, will limit the nutrients entering the lake, starving the plants and algae of the food they need for growth.



9353 Hill Road • Swartz Creek, MI 48473
 (810) 635-4400 • Fax (810) 635-4404


www.lakeproinc.com



Transparency (feet)					
	April	May	June	July	August
West	2.9	7.7	8.8	9.8	6.9
East	3.1	9.3	11.5	11.5	6.4
Season Average					7.8
Trophic State Index					49

Target Range: > 6.5 feet

Less chlorophyll generally leads to higher transparency. Despite the downward trend of chlorophyll, the transparency also trended down. In 2014, additional tests revealed the brown color in the early summer was due to high concentrations of the algae Planktothrix. The decreasing transparency is a concern for the lake. It ruins the appearance of the lake and can pose a threat to the safety of the lake, because swimmers cannot accurately judge depth or see underwater obstructions.



pH (Standard Units)					
	April	May	June	July	August
West	8.3	8.4	8.3	8.0	7.5
East	8.3	8.8	7.9	8.0	7.8
Season Average					8.1


Target Range: 7.0 – 9.0 S.U.

The pH at the lake surface trended slightly downward over the testing history. It is important to look for the trend to flatten with additional years of data. Some fluctuation is expected, but major shifts or changes beyond the target range may indicate a larger problem.



9353 Hill Road • Swartz Creek, MI 48473
 (810) 635-4400 • Fax (810) 635-4404


www.lakeproinc.com



Total Dissolved Solids (ppm)					
	April	May	June	July	August
West	391	390	369	365	355
East	390	389	368	364	359
Season Average					374

Target Range: 0 – 1,000 ppm

The total dissolved solids increased slowly during the testing history, showing that the lake is accumulating more substances. Continuing to harvest will remove substances from the lake, including nutrients, as they are bound into the plant biomass. Fewer plants also allows easier and faster water movement, which prevents deposition of sediments and increased flushing of excess substances.



Conductivity (µS)					
	April	May	June	July	August
West	782	779	739	729	712
East	781	779	735	728	718
Season Average					748


Target Range: 0 – 1,500 µS

Like the TDS, Conductivity increased over the testing history. Conductivity is a function of TDS and measures the amount of ionic molecules in the water (which conduct electricity; usually salts). We will look for this trend to slow in future years of testing.



9353 Hill Road • Swartz Creek, MI 48473
 (810) 635-4400 • Fax (810) 635-4404


www.lakeproinc.com



Alkalinity (ppm)					
	April	May	June	July	August
West	145	141	130	117	105
East	151	148	138	127	113
Season Average					132

Target Range: 0 – 250 ppm

Alkalinity was first tested in 2012, so the historical data is limited. During the past seven years, alkalinity slowly increased. Additional carbonates are beneficial to the lake because they are natural buffers. Also, the increase of carbonates contributed to the increases in total dissolved solids and conductivity.



Sulfate (ppm)					
	April	May	June	July	August
West	12.9	12.9	13.6	13.0	13.3
East	12.6	13.6	12.9	13.1	14.2
Season Average					13.2


Target Range: 3 – 30 ppm

Sulfate has not changed significantly since 2002. It is important that this parameter stay within the target range and any significant increases will be investigated.



9353 Hill Road • Swartz Creek, MI 48473
 (810) 635-4400 • Fax (810) 635-4404


www.lakeproinc.com



Fluoride (ppm)					
	April	May	June	July	August
West	0.09	0.10	0.08	0.08	0.10
East	0.08	0.09	0.10	0.08	0.10
Season Average					0.09

Target Range: 0.01 – 0.30 ppm

Fluoride decreased in the lake since testing began. It is important that this parameter stay within the target range and any significant increases will be investigated.



Chloride (ppm)					
	April	May	June	July	August
West	133	128	121	119	109
East	138	135	129	126	120
Season Average					126

Target Range: 0 – 230 ppm

The trend line for Chloride was flat, but recent years were below the trend. With concerns about road salt entering lakes, it is important that this parameter remain within the target range and any increases will be investigated.



9353 Hill Road • Swartz Creek, MI 48473
(810) 635-4400 • Fax (810) 635-4404

www.lakeproinc.com

Analysis Information

Temperature:	The water temperature directly affects the amount of oxygen that can dissolve into the water. The temperature of surface waters is not indicative of the entire water column.
Dissolved Oxygen:	D.O. is a measure of the amount of oxygen dissolved in the water. This oxygen is available to fish and other animals for respiration. Vegetation generally increases DO, particularly during the day and early evening. Animals and other respiring organisms consume the oxygen, mostly during the day. Oxygen is also added to the lake through wave action, rain, fountains and aerators.
Total Phosphorus:	Phosphorus is an essential nutrient for plant growth. However, concentrations exceeding 100 ppb can impair the water and results in nuisance vegetation growth.
Phosphates:	Phosphate is the form of phosphorous that is most readily available to plants and algae.
Nitrate:	Nitrogen is also essential for plant growth. Nitrate is the predominant form of nitrogen in water. Excessive nitrate concentrations may also result in pollution and increased vegetation.
Chlorophyll-a:	Chlorophyll-a is a direct measurement of the amount of green pigment produced by plants and phytoplankton. This indicates the amount of plant growth and is used to calculate a Trophic State Index.
Transparency:	The ability of light to penetrate the water column is determined by the amount of dissolved and suspended particles in the water. Although aesthetically desirable, transparent water allows increased light to reach the lake bed and may result in vegetation growth.
pH:	pH is a measure of acidity or alkalinity. pH is a general measure of lake health and can roughly indicate the range of other measurements such as alkalinity and hardness.
TDS:	Total Dissolved Solids is the amount of all organic and inorganic substances in the water in a molecular or ionized state. Higher values generally indicate richer and more productive water. Lower values usually indicate cleaner and less productive water.
Conductivity:	Conductivity is a measure of the ability of water to conduct electricity. Dissolved ions in the water increase conductivity, thus TDS and Conductivity are closely related.
Alkalinity:	Alkalinity refers to the ability of the water to neutralize acids, mainly through the hydrogenation of carbonate ions. Therefore, the alkalinity is expressed as "ppm as CaCO ₃ ". However, other basic molecules in the water can also contribute to alkalinity.
Sulfate:	Sulfate occurs naturally as minerals, such as calcium sulfate and magnesium sulfate. In fresh water, sulfate is usually the second or third most abundant anion. Other sources of sulfate include water material from pulp mills, steel mills, food processing operations, and municipal wastes. Under low oxygen conditions, sulfate can be reduced to hydrogen sulfide gas, which smells like rotten eggs.
Fluoride:	Fluoride may occur naturally or be added to public drinking water supplies.
Chloride:	Chloride is one of the major anions found in water and sewage. The presence of chlorides may be due to water passing through salt formations in the earth or pollution from industrial processes, domestic wastes, or road salt. The salt content of water affects the distribution of plant and animal life in an aquatic system, based on the amount of salt they can tolerate.



9353 Hill Road • Swartz Creek, MI 48473
(810) 635-4400 • Fax (810) 635-4404

www.lakeproinc.com

Trophic States

- Oligotrophic:** Water is very clear. Nutrient levels are generally low. Plant and algae productivity are also low. Sufficient dissolved oxygen in the bottom, cooler waters allows cold-water fish to survive, such as salmon and trout.
- Mesotrophic:** Water is moderately clear. Nutrient levels are slightly elevated. Plant and algae productivity are present, but generally not a nuisance. Oxygen and temperature in the lower portion of the lake allow walleye and perch to survive.
- Eutrophic:** Water is not clear due to high nutrients levels, increased turbidity, and excessive algal growth. There is no oxygen in the bottom, cooler waters, restricting the lake to warm water species, such as bass and bluegill.
- Hypereutrophic:** Nutrient levels are extremely high, promoting very high algae productivity. Blue-green algae blooms are likely. High turbidity and algae growth make the water opaque. Little plant growth is restricted to invasive plants. The only fish that can survive this environment are rough fish, such as carp, catfish, and mudminnows.

Sample Sites:

